Journal of Immunological Methods, 134 (1990) 283–284
Elsevier

Contents to volume 134 (1990)

Number 1 – 6 November 1990

Announcement from the publisher v

Review article

Detection of peptide-MHC class II complexes on the surface of intact cells
R. Busch and J.B. Rothbard (U.S.A.) 1

Research reports

Multi-pin peptide synthesis strategy for T cell determinant analysis
N.J. Maeji, A.M. Bray and H.M. Geysen (Australia) 23

Development of microfusion techniques to generate human hybridomas
S. Foung, S. Perkins, K. Kafadar, P. Gessner and U. Zimmermann (U.S.A., F.R.G.) 35

Efficient hybridization of mouse-human cell lines by means of hypo-osmolar electrofusion
U. Zimmermann, P. Gessner, R. Schnettler, S. Perkins and S.K.H. Foung (F.R.G., U.S.A.) 43

Methods to quantitate human haptoglobin by complexation with hemoglobin
S. Cerda and S.K. Oh (U.S.A.) 51

Cytotoxicity assays using cryopreserved target cells pre-labeled with the fluorescent marker europium
D.T. Maley and P. Simon (U.S.A.) 61

A silicon sensor-based filtration immunoassay using biotin-mediated capture
J.D. Olson, P.R. Panfili, R. Armenta, M.B. Femmel, H. Merrick, J. Gumperz, M. Goltz and R.F. Zuk (U.S.A.) 71

A rapid method for monitoring DNA labelling reactions with haptens
A. Corti, P. Tripputi and G. Cassani (Italy) 81

Monoclonal antibodies distinguishing α and β forms of calcitonin gene-related peptide
D.P. Andrew, T.D. Bidgood, C. Bose, D. Brown, G. Galfre and M. Sherwood (U.K.) 87

Comparison of an enzyme-linked immunosorbent assay (ELISA) with a radioimmunoassay (RIA) for the measurement of rat insulin
H.V. Webster, A.J. Bone, K.A. Webster and T.J. Wilkin (U.K.) 95

Quantification of acute phase proteins in rat serum and in the supernatants of a cultured rat hepatoma cell line and cultured primary hepatocytes by an enzyme-linked immunosorbent assay
M. Ikawa and Y. Shozen (Japan) 101

Affinity immunoblotting detection of serum monoclonal immunoglobulins reactive with glycosphingolipids
M.O. Jauberteau, J.M. Cook, M. Drouet and J.L. Preud'homme (France) 107

Quantitative determination of retrovirus-specific immune complexes
H. Schetters, H. Rohmer, R. Hehlmann and V. Erfle (F.R.G.) 113

Murine monoclonal anti-idiotype antibody (α) as a probe to detect human monoclonal antibody bound to human tumor tissues
H. Saito, D.-R. Wen, S. Yamamoto, T. Yamamoto, R.E. Saxton, A.J. Cochran and R.F. Irie (U.S.A.) 121

Use of a monoclonal antibody specific for a protein epitope of *Echinococcus granulosus* antigen 5 in a competitive antibody radioimmunoassay for diagnosis of hydatid disease
M. Chamekh, B. Facon, C. Dissous, A. Haque and A. Capron (France) 129

Development of technology for linking photosensitizers to a model monoclonal antibody
F.N. Jiang, S. Jiang, D. Liu, A. Richter and J.G. Levy (Canada, China) 139

Notes to authors 151

Number 2 – 5 December 1990

Announcement from the publisher iii

Research reports

A new method for removal of mononuclear phagocytes from heterogeneous cell populations in vitro, using the liposome-mediated macrophage 'suicide' technique
I. Claassen, N. Van Rooijen and E. Claassen (The Netherlands) 153

Quantification of α-fetoprotein and transferrin endocytosis by lymphoid cells using flow cytometry
J.M. Torres, C. Esteban, J. Aguilar, Z. Mishal and J. Uriel . (France) 163
Particle counting immunoassay of choriogonadotropin using monoclonal antibodies
M. Halfiants, N. Tasiaux, L. Van Krieken, R. De Hertogh and D. Collet-Cassart (Belgium) 171
Comparison of various methods for monitoring hybridoma cell proliferation
D. Duval, C. Demangel, I. Geahel, K. Blondeau and A. Marcadet (France) 177
Subnucleosome structures as substrates in enzyme-linked immunosorbent assays
R.W. Burlingame and R.L. Rubin (U.S.A.) 187
A new assay for cell death
I.P. Beletsky and S.R. Umansky (U.S.S.R.) 201
Antibody-coated liposomes as a particulate solid phase for immunoassays. Measurement of urinary 'micro-albumin'
S.J. Frost, G.B. Firth and J. Chakraborty (U.K.) 207
Comparison of methods for the analysis of outer membrane antigens of *Neisseria meningitidis* by Western blotting
R.L. Davies, R.A. Wall and S.P. Borriello (U.K.) 215
The use of *N*-(aminobenzoyloxy)succinimide as a two-level heterobifunctional agent for the preparation of hapten-protein conjugates. Daunomycin as a model hapten with an amino group
K. Fujiwara, N. Matsumoto, T. Kitagawa and K. Inouye (Japan) 227
A new one-step method for the isolation of human mononuclear cells
T.C. Ford and D. Rickwood (U.K.) 237
Detection and quantitation of HLA class II molecules on keratinocytes by quantitative immunofluorescence
S.R. Kinn, A. Curtis and J.A.T. Dow (U.K.) 243
A rapid and sensitive immunoassay for tumor necrosis factor using magnetic monodisperse polymer particles
N.-B. Liabakk, K. Nustad and T. Espevik (Norway) 253
Competitive ELISA for the measurement of tau protein in Alzheimer's disease
C.R. Harrington, P.C. Edwards and C.M. Wischik (U.K.) 261
Immunoblotting techniques for the detection of low level homogeneous immunoglobulin components in serum
F.J.M. Nooij, A.J. Van der Sluijs-Gelling, C.M. Jol-Van der Zijde, M.J.D. Van Tol, H. Haas and J. Radl (The Netherlands, F.R.G.) 273

Contents to volume 134 (1990) 283

Notes to authors 285